

## NEWHATCH-C2-M® Newcastle-Bronchitis Vaccine

B1 Type, C2 Strain, Mass. Type, Live Virus

### Product Information

**NEWHATCH-C2-M®** combines the patented, virtually non-reactive C2 strain of B1 type Newcastle Disease (ND) virus with the well-established infectious bronchitis (IB) protection provided by the Intervet mild Massachusetts strain of IB. It is a lyophilized vaccine approved for spray vaccination of chickens one day of age or older for protection against Newcastle disease and infectious bronchitis disease (Massachusetts type).



### NEWHATCH-C2-M® contains:

- Patented C2 strain of B1 type Newcastle Disease virus.
- Mild Massachusetts type of infectious bronchitis virus.

### Product Features:

- C2 strain of B1 type Newcastle minimizes reaction to day-of-age respiratory vaccination in broiler chicks.
- NEWHATCH-C2-M® eliminates problems with lingering hatchery reaction prior to field boost.
- Improved livability and performance due to minimized reactions.
- The virtually non-reactive C2 strain permits the use of a full dose of vaccine for better protection against both ND and IB.
- Approved for hatchery application.

## Description:

**Newhatch-C2-M®** is a lyophilized vaccine prepared from specific pathogen free (SPF) substrates and contains the patented C2 strain of B1 Type Newcastle disease virus and the mild Massachusetts type of infectious bronchitis virus.

## Indications for Use:

Newhatch-C2-M® is indicated for the vaccination of healthy chickens at one day of age or older by coarse spray for protection against Newcastle disease and infectious bronchitis (Massachusetts type).

## Dosage and Administration:

The time of administration should be determined by the degree of Newcastle disease and infectious bronchitis exposure, type of chicken being vaccinated, and the current management procedures. **Newhatch-C2-M®** is approved for use via coarse spray at one day of age or older.

## Vaccination:

The best protection is obtained when the vaccinated chickens have time to develop immunity prior to exposure to Newcastle disease or infectious bronchitis virus. Good management practices are recommended to reduce the exposure to these viruses for at least three weeks following vaccination.

## Cautions:

1. Avoid exposure of this vaccine to sunlight.
2. Vaccinate only healthy birds.
3. All susceptible birds on the same premises should be vaccinated at the same time.
4. Efforts should be taken to reduce stress conditions at the time of vaccination and during the reaction period.
5. Do not dilute the vaccine or otherwise stretch the dosage of the vaccine.
6. Use entire contents of the vial when first opened.
7. Do not vaccinate chickens within 21 days of slaughter.


## How Supplied

10 x 10,000 dose vials

## Product Code

NB-208-10

## Cap Color Code



U.S. Patent No. 5,750,111